

The Ergodic Hierarchy of Mixing, van der Corput's difference theorem, and the ergodic theory of noncommuting operators

Nilpotent structures in topological dynamics, ergodic theory and combinatorics at Pałac Będlewo

Based on <https://arxiv.org/abs/2303.11832>

Sohail Farhangi
Slides available on sohailfarhangi.com

June 5, 2023

- 1 Review
 - Van der Corput's difference theorem and some applications
 - Mixing properties
- 2 Mixing van der Corput difference theorems
- 3 Ergodic theorems for noncommuting operators
 - Background
 - New results from mixing vdCs
- 4 Examples of systems with singular spectrum

Table of Contents

1 Review

- Van der Corput's difference theorem and some applications
- Mixing properties

2 Mixing van der Corput difference theorems

3 Ergodic theorems for noncommuting operators

- Background
- New results from mixing vdCs

4 Examples of systems with singular spectrum

The Classical van der Corput Difference Theorem

Definition

A sequence $(x_n)_{n=1}^{\infty} \subseteq [0, 1]$ is **uniformly distributed** if for any open interval $(a, b) \subseteq [0, 1]$ we have

$$\lim_{N \rightarrow \infty} \frac{1}{N} |\{1 \leq n \leq N \mid x_n \in (a, b)\}| = b - a. \quad (1)$$

Theorem (van der Corput, [26])

If $(x_n)_{n=1}^{\infty} \subseteq [0, 1]$ is such that $(x_{n+h} - x_n)_{n=1}^{\infty}$ is uniformly distributed for every $h \in \mathbb{N}$, then $(x_n)_{n=1}^{\infty}$ is itself uniformly distributed.

Corollary

If $\alpha \in \mathbb{R}$ is irrational, then $(n^2\alpha)_{n=1}^{\infty}$ is uniformly distributed.

Theorem (HvdCDT1, Bergelson [3, Theorem 1.4])

If \mathcal{H} is a Hilbert space and $(x_n)_{n=1}^{\infty} \subseteq \mathcal{H}$ is a bounded sequence satisfying

$$\lim_{N \rightarrow \infty} \frac{1}{N} \sum_{n=1}^N \langle x_{n+h}, x_n \rangle = 0, \quad (2)$$

for every $h \in \mathbb{N}$, then

$$\lim_{N \rightarrow \infty} \left\| \frac{1}{N} \sum_{n=1}^N x_n \right\| = 0. \quad (3)$$

Theorem (HvdCDT2, Bergelson [3, Page 3])

If \mathcal{H} is a Hilbert space and $(x_n)_{n=1}^{\infty} \subseteq \mathcal{H}$ is a bounded sequence satisfying

$$\lim_{h \rightarrow \infty} \limsup_{N \rightarrow \infty} \left| \frac{1}{N} \sum_{n=1}^N \langle x_{n+h}, x_n \rangle \right| = 0, \text{ then} \quad (4)$$

$$\lim_{N \rightarrow \infty} \left\| \frac{1}{N} \sum_{n=1}^N x_n \right\| = 0. \quad (5)$$

Theorem (HvdCDT3, Bergelson [3, Theorem 1.5])

If \mathcal{H} is a Hilbert space and $(x_n)_{n=1}^{\infty} \subseteq \mathcal{H}$ is a bounded sequence satisfying

$$\lim_{H \rightarrow \infty} \frac{1}{H} \sum_{h=1}^H \limsup_{N \rightarrow \infty} \left| \frac{1}{N} \sum_{n=1}^N \langle x_{n+h}, x_n \rangle \right| = 0, \text{ then} \quad (6)$$

$$\lim_{N \rightarrow \infty} \left\| \frac{1}{N} \sum_{n=1}^N x_n \right\| = 0. \quad (7)$$

Question

Why would we ever use HvdCDT1 or HvdCDT2 when they are both corollaries of HvdCDT3? Why are there at least 3 Hilbertian vdCDTs and only 1 vdCDT in the theory of uniform distribution?

Applications of HvdCDTs 1/2

Theorem (Poincaré)

For any measure preserving system (m.p.s.) (X, \mathcal{B}, μ, T) , and any $A \in \mathcal{B}$ with $\mu(A) > 0$, there exists $n \in \mathbb{N}$ for which

$$\mu(A \cap T^{-n}A) > 0. \quad (8)$$

Does not need vdCDT.

Theorem (Furstenberg-Sárközy)

For any m.p.s. (X, \mathcal{B}, μ, T) , and any $A \in \mathcal{B}$ with $\mu(A) > 0$, there exists $n \in \mathbb{N}$ for which

$$\mu(A \cap T^{-n^2}A) > 0. \quad (9)$$

Furstenberg's proof in [17, Proposition 1.3] uses a form of vdCDT since it uses the uniform distribution of $(n^2\alpha)_{n=1}^{\infty}$. See also [4, Theorem 2.1] for a proof using HvdCDT1 directly.

Applications of HvdCDTs 2/2

Theorem (Furstenberg, [17])

For any m.p.s. (X, \mathcal{B}, μ, T) , any $A \in \mathcal{B}$ with $\mu(A) > 0$, and any $\ell \in \mathbb{N}$, there exists $n \in \mathbb{N}$ for which

$$\mu(A \cap T^{-n}A \cap T^{-2n}A \cap \cdots \cap T^{-\ell n}A) > 0. \quad (10)$$

The proof presented in [9] uses HvdCT3 as Theorem 7.11, and the proof in [18] uses a variation.

Theorem (Bergelson and Leibman, [6])

For any m.p.s. $(X, \mathcal{B}, \mu, \{T_i\}_{i=1}^\ell)$ with the T_i s commuting, any $A \in \mathcal{B}$ with $\mu(A) > 0$, and any $\{p_i(x)\}_{i=1}^\ell \subseteq x\mathbb{N}[x]$, there exists $n \in \mathbb{N}$ for which

$$\mu(A \cap T_1^{-p_1(n)}A \cap T_2^{-p_2(n)}A \cap \cdots \cap T_\ell^{-p_\ell(n)}A) > 0. \quad (11)$$

Uses an equivalent form of HvdCT3 as Lemma 2.4.

Some of the Ergodic Hierarchy of Mixing

Definition

Let $\mathcal{X} = (X, \mathcal{B}, \mu, T)$ be a m.p.s. If for every $f, g \in L_0^2(X, \mu)$

- ① $\lim_{N \rightarrow \infty} \frac{1}{N} \sum_{n=1}^N \langle U_T^n f, g \rangle = 0$, then \mathcal{X} is **ergodic**.
- ② $\lim_{N \rightarrow \infty} \frac{1}{N} \sum_{n=1}^N |\langle U_T^n f, g \rangle| = 0$, then \mathcal{X} is **weakly mixing**,
- ③ $\lim_{n \rightarrow \infty} \langle U_T^n f, g \rangle = 0$, then \mathcal{X} is **strongly mixing**,
- ④ and if $L_0^2(X, \mu)$ has an orthogonal basis of the form $\{U_T^n f_m\}_{n,m \in \mathbb{Z}}$, then \mathcal{X} has **Lebesgue spectrum**.
- ⑤ which is the same as $(\langle U_T^n f, g \rangle)_{n=1}^\infty$ being Fourier coefficients of some $h \in L^1([0, 1], \mathcal{L})$, where \mathcal{L} is the Lebesgue measure.

These definitions also apply to individual elements $f \in L_0^2(X, \mu)$.

The Symmetric Ergodic Hierarchy of Mixing

Theorem

Let $\mathcal{X} = (X, \mathcal{B}, \mu, T)$ be a m.p.s. If for every $f \in L_0^2(X, \mu)$

- ① $\lim_{N \rightarrow \infty} \frac{1}{N} \sum_{n=1}^N \langle U_T^n f, f \rangle = 0$, then \mathcal{X} is **ergodic**,
- ② $\lim_{N \rightarrow \infty} \frac{1}{N} \sum_{n=1}^N |\langle U_T^n f, f \rangle| = 0$, then \mathcal{X} is **weakly mixing**,
- ③ $\lim_{n \rightarrow \infty} \langle U_T^n f, f \rangle = 0$, then \mathcal{X} is **strongly mixing**,
- ④ \mathcal{X} has **Lebesgue spectrum** if $(\langle U_T^n f, f \rangle)_{n=1}^\infty$ are the Fourier coefficients of some $h \in L^1([0, 1], \mathcal{L})$ taking nonnegative real values.

This theorem also applies to individual elements $f \in L_0^2(X, \mu)$.

Table of Contents

1 Review

- Van der Corput's difference theorem and some applications
- Mixing properties

2 Mixing van der Corput difference theorems

3 Ergodic theorems for noncommuting operators

- Background
- New results from mixing vdCs

4 Examples of systems with singular spectrum

Theorem (F. 2022)

If $(x_n)_{n=1}^{\infty} \subseteq \mathcal{H}$ is a bounded sequence satisfying

$$\lim_{H \rightarrow \infty} \frac{1}{H} \sum_{h=1}^H \limsup_{N \rightarrow \infty} \left| \frac{1}{N} \sum_{n=1}^N \langle x_{n+h}, x_n \rangle \right| = 0, \quad (12)$$

then $(x_n)_{n=1}^{\infty}$ is a *nearly weakly mixing sequence*. This means that for any other bounded sequence $(y_n)_{n=1}^{\infty} \subseteq \mathcal{H}$ we morally (*but not literally*) have that

$$\lim_{H \rightarrow \infty} \frac{1}{H} \sum_{h=1}^H \left| \lim_{N \rightarrow \infty} \frac{1}{N} \sum_{n=1}^N \langle x_{n+h}, y_n \rangle \right| = 0. \quad (13)$$

Loosely speaking, this can be interpreted as a *weak mixing* in any ultrapower \mathcal{H} of \mathcal{H} with respect to a unitary operator induced by the left shift. Note that elements of \mathcal{H} are sequences in \mathcal{H} .

A Lebesgue spectrum vdCdt

Theorem (F. 2023)

If $(x_n)_{n=1}^{\infty} \subseteq \mathcal{H}$ is a bounded sequence satisfying for all $h \in \mathbb{N}$

$$\sum_{h=1}^{\infty} \limsup_{N \rightarrow \infty} \left| \frac{1}{N} \sum_{n=1}^N \langle x_{n+h}, x_n \rangle \right|^2 < \infty, \quad (14)$$

then $(x_n)_{n=1}^{\infty}$ is a **spectrally Lebesgue sequence**. If $\mathcal{H} = L^2(X, \mu)$ and $(y_n)_{n=1}^{\infty} \subseteq L^{\infty}(X, \mu)$ is bounded and **spectrally singular**, then

$$\lim_{N \rightarrow \infty} \left\| \frac{1}{N} \sum_{n=1}^N x_n y_n \right\| = 0. \quad (\#)$$

Upgrading from weak convergence to the strong convergence in **#** necessitates a new proof of the classical vdCDT. See [10, Chapter 2] for variations of MvdCT related to other levels of mixing, as well as uniform distribution. See also [25].

Table of Contents

1 Review

- Van der Corput's difference theorem and some applications
- Mixing properties

2 Mixing van der Corput difference theorems

3 Ergodic theorems for noncommuting operators

- Background
- New results from mixing vdCs

4 Examples of systems with singular spectrum

Failure of noncommutative ergodic theorems

Theorem (Frantzkinakis, Lesigne, Wierdl [15, Lemma 4.1])

Let $a, b : \mathbb{N} \rightarrow \mathbb{Z} \setminus \{0\}$ be injective sequences and F be any subset of \mathbb{N} . Then there exist a probability space (X, \mathcal{B}, μ) , measure preserving automorphisms $T, S : X \rightarrow X$, both of them Bernoulli, and $A \in \mathcal{B}$, such that

$$\mu(T^{-a(n)}A \cap S^{-b(n)}A) = \begin{cases} 0 & \text{if } n \in F, \\ \frac{1}{4} & \text{if } n \notin F. \end{cases} \quad (15)$$

See also Furstenberg [18, Page 40], Berend [2, Example 7.1], as well as Bergelson and Leibman [7].

Noncommutative ergodic theorems 1/2

Theorem (Frantzkinakis [12, Corollary 1.7])

Let $a : \mathbb{R}_+ \rightarrow \mathbb{R}$ be a Hardy field function for which there exist some $\epsilon > 0$ and $d \in \mathbb{Z}_+$ satisfying

$$\lim_{t \rightarrow \infty} \frac{a(t)}{t^{d+\epsilon}} = \lim_{t \rightarrow \infty} \frac{t^{d+1}}{a(t)} = \infty. \quad (\text{e.g. } a(t) = t^{1.5}) \quad (16)$$

Furthermore, let (X, \mathcal{B}, μ) be a probability space and $T, S : X \rightarrow X$ be measure preserving transformations. Suppose that the system (X, \mathcal{B}, μ, T) has **zero entropy**. Then

(i) For every $f, g \in L^\infty(X, \mu)$ we have

$$\lim_{N \rightarrow \infty} \frac{1}{N} \sum_{n=1}^N T^n f \cdot S^{\lfloor a(n) \rfloor} g = \mathbb{E}[f | \mathcal{I}_T] \cdot \mathbb{E}[g | \mathcal{I}_S], \quad (17)$$

where the limit is taken in $L^2(X, \mu)$.

Theorem (Continued)

(ii) For every $A \in \mathcal{B}$ we have

$$\lim_{N \rightarrow \infty} \frac{1}{N} \sum_{n=1}^N \mu(A \cap T^{-n}A \cap S^{-\lfloor a(n) \rfloor}A) \geq \mu(A)^3. \quad (18)$$

Frantzikinakis and Host [13] a similar theorem is proven for $a(n) = p(n)$ with $p(x) \in \mathbb{Z}[x]$ of degree at least 2. Rohlin [23] showed that every T with **singular spectrum** must also have **zero entropy**. Note that the Horocycle flow has **zero entropy** [19] and **Lebesgue spectrum** [22]

Theorem (F., 2023)

Let (X, \mathcal{B}, μ) be a probability space and let $T, S : X \rightarrow X$ be measure preserving automorphisms for which T has **singular spectrum**. Let $(k_n)_{n=1}^{\infty} \subseteq \mathbb{N}$ be a sequence for which $((k_{n+h} - k_n)\alpha)_{n=1}^{\infty}$ is uniformly distributed in the orbit closure of α for all $\alpha \in \mathbb{R}$ and $h \in \mathbb{N}$.

① For any $f, g \in L^{\infty}(X, \mu)$ we have

$$\lim_{N \rightarrow \infty} \frac{1}{N} \sum_{n=1}^N T^n f \cdot S^{k_n} g = \mathbb{E}[f | \mathcal{I}_T] \mathbb{E}[g | \mathcal{I}_S], \quad (19)$$

with convergence taking place in $L^2(X, \mu)$.

Theorem (Continued)

(ii) *If $A \in \mathcal{B}$ then*

$$\lim_{N \rightarrow \infty} \frac{1}{N} \sum_{n=1}^N \mu(A \cap T^{-n}A \cap S^{-k_n}A) \geq \mu(A)^3. \quad (20)$$

(iii) *If we only assume that $((k_{n+h} - k_n)\alpha)_{n=1}^{\infty}$ is uniformly distributed for all $\alpha \in \mathbb{R} \setminus \mathbb{Q}$ and $h \in \mathbb{N}$, then (i) and (ii) hold when S is **totally ergodic**.*

Examples include $k_n = \lfloor a(n) \rfloor$ with $a(n)$ being as in frame 19, $k_n = \lfloor n^2 \log^2(n) \rfloor$, and for part (iii) we may take $k_n = p(n)$ for $p(x) \in x\mathbb{Z}[x]$ with degree at least 2.

Application 2/4 (A special case)

Theorem (F., 2023)

Let (X, \mathcal{B}, μ) be a probability space and $T, S : X \rightarrow X$ be measure preserving automorphisms. Suppose that T has **singular spectrum** and S is **totally ergodic**. Let $p_1, \dots, p_K \in \mathbb{Q}[x]$ be integer polynomials for which $\deg(p_1) \geq 2$ and $\deg(p_i) \geq 2 + \deg(p_{i-1})$. For any $f, g_1, \dots, g_K \in L^\infty(X, \mu)$, we have

$$\lim_{N \rightarrow \infty} \frac{1}{N} \sum_{n=1}^N T^n f \prod_{i=1}^K S^{p_i(n)} g_i = \mathbb{E}[f | \mathcal{I}_T] \prod_{i=1}^K \int_X g_i d\mu, \quad (21)$$

with convergence taking place in $L^2(X, \mu)$.

Application 3/4 (A special case)

Theorem (F., 2023)

Let (X, \mathcal{B}, μ) be a probability space and $T, R, S : X \rightarrow X$ be measure preserving automorphisms. Suppose that T has **singular spectrum**, R and S commute, and S is **weakly mixing**. Let $\ell \in \mathbb{N}$ and let $p_1, \dots, p_\ell \in \mathbb{Q}[x]$ be pairwise essentially distinct integer polynomials, each having degree at least 2. For any $f, h, g_1, \dots, g_\ell \in L^\infty(X, \mu)$ satisfying $\int_X g_j d\mu = 0$ for some $1 \leq j \leq \ell$, we have

$$\lim_{N \rightarrow \infty} \frac{1}{N} \sum_{n=1}^N T^n f \cdot R^n h \cdot \prod_{j=1}^{\ell} S^{p_j(n)} g_j = 0, \quad (22)$$

with convergence taking place in $L^2(X, \mu)$.

An example

Consider the m.p.s. $([0, 1]^2, \mathcal{B}, \mathcal{L}^2, T, S)$ with $S(x, y) = (x + 2\alpha, y + x)$ for some $\alpha \in \mathbb{R} \setminus \mathbb{Q}$, and $T(x, y) = (x, y + x)$. We see that $([0, 1]^2, \mathcal{B}, \mathcal{L}^2, S)$ and $([0, 1]^2, \mathcal{B}, \mathcal{L}^2, T)$ are both **zero entropy** systems that are not **weakly mixing**, and the former is **totally ergodic**. Furthermore, T and S generate a 2-step nilpotent group. For $f_0(x, y) = e^{2\pi i(x-y)}$, $f_1(x, y) = e^{2\pi iy}$, and $f_2(x, y) = e^{-2\pi ix}$, we see that

$$\begin{aligned} & \lim_{N \rightarrow \infty} \frac{1}{N} \sum_{n=1}^N T^n f_0(x, y) S^n f_1(x, y) S^{\frac{1}{2}(n^2-n)} f_2(x, y) \\ &= \lim_{N \rightarrow \infty} \frac{1}{N} \sum_{n=1}^N e^{2\pi i \left((1-n)x - y + y + nx + (n^2-n)\alpha - x - (n^2-n)\alpha \right)} = 1 \neq 0. \end{aligned}$$

Sets of K but not $K + 1$ recurrence?

Theorem (Frantzkinakis, Lesigne, Wierdl [14])

Let $k \geq 2$ be an integer and $\alpha \in \mathbb{R}$ be irrational. Let

$$R_k = \left\{ n \in \mathbb{N} \mid n^k \alpha \in \left[\frac{1}{4}, \frac{3}{4} \right] \right\}.$$

(i) If (X, \mathcal{B}, μ) is a probability space and

$S_1, S_2, \dots, S_{k-1} : X \rightarrow X$ are commuting measure preserving transformations, then for any $A \in \mathcal{B}$ with $\mu(A) > 0$, there exists $n \in R_k$ for which

$$\mu(A \cap S_1^{-n}A \cap S_2^{-n}A \cap \dots \cap S_{k-1}^{-n}A) > 0. \quad (23)$$

(ii) There exists a m.p.s. (X, \mathcal{B}, μ, T) and a set $A \in \mathcal{B}$ satisfying $\mu(A) > 0$ such that for all $n \in R_k$ we have

$$\mu(A \cap T^{-n}A \cap T^{-2n}A \cap \dots \cap T^{-kn}A) = 0. \quad (24)$$

Application 4/4

Theorem (F., 2023)

Let $k \geq 2$ be an integer and $\alpha \in \mathbb{R}$ be irrational. Let $R_k = \{n \in \mathbb{N} \mid n^k \alpha \in [\frac{1}{4}, \frac{3}{4}]\}$. Let (X, \mathcal{B}, μ) be a probability space and $S_1, S_2, \dots, S_{k-1} : X \rightarrow X$ commuting measure preserving automorphisms. Let $T : X \rightarrow X$ be a measure preserving automorphism with **singular spectrum**, and for which $\{T, S_1, S_2, \dots, S_{k-1}\}$ generate a nilpotent group. For any $A \in \mathcal{B}$ with $\mu(A) > 0$, there exists $n \in R$ for which

$$\mu(A \cap T^{-n}A \cap S_1^{-n}A \cap S_2^{-n}A \cap \dots \cap S_{k-1}^{-n}A) > 0. \quad (25)$$

Since the system $(\mathbb{T}^2, \mathcal{B}^2, \mathcal{L}^2, T)$ with $T(x, y) = (x + \alpha, y + x)$ can be used in item (ii) of the last slide when $k = 2$, the current theorem does not hold for a general T with 0 entropy. Also note that the maximal spectral type of T is $\mathcal{L} + \sum_{n \in \mathbb{Z}} \delta_{n\alpha}$.

Table of Contents

1 Review

- Van der Corput's difference theorem and some applications
- Mixing properties

2 Mixing van der Corput difference theorems

3 Ergodic theorems for noncommuting operators

- Background
- New results from mixing vdCs

4 Examples of systems with singular spectrum

Examples of systems with singular spectrum

In [5, Proposition 2.9] it is shown that if (X, \mathcal{B}, μ) is a standard probability space, and $\text{Aut}(X, \mathcal{B}, \mu)$ is endowed with the strong operator topology, then the set of transformations that are **weakly mixing** and rigid is a generic set. Since any rigid automorphism has **singular spectrum**, we see that the set of singular automorphisms is generic. Now let $\mathcal{S} \subseteq \text{Aut}(X, \mathcal{B}, \mu)$ denote the collection of **strongly mixing** transformation, and note that \mathcal{S} is a meager set since an automorphism cannot simultaneously be rigid and **strongly mixing**. Since \mathcal{S} is not a complete metric space with respect to the topology induced by the strong operator topology, a new topology was introduced in [24], with respect to which \mathcal{S} is a complete metric space. It is shown in the Corollary to Theorem 7 of [24] that a generic $T \in \mathcal{S}$ has **singular spectrum**, and such a T is mixing of all orders due a well known result of Host [20]. See [11] and [21] for concrete examples of $T \in \mathcal{S}$ that have **singular spectrum**. See also [1], [8], and [16].

References |

- [1] C. Aistleitner and M. Hofer.
On the maximal spectral type of a class of rank one transformations.
Dyn. Syst., 27(4):515–523, 2012.
- [2] D. Berend.
Joint ergodicity and mixing.
J. Analyse Math., 45:255–284, 1985.
- [3] V. Bergelson.
Weakly mixing PET.
Ergodic Theory Dynam. Systems, 7(3):337–349, 1987.

References II

- [4] V. Bergelson.
Ergodic Ramsey theory—an update.
In *Ergodic theory of \mathbb{Z}^d actions (Warwick, 1993–1994)*, volume 228 of *London Math. Soc. Lecture Note Ser.*, pages 1–61. Cambridge Univ. Press, Cambridge, 1996.
- [5] V. Bergelson, A. del Junco, M. Lemańczyk, and J. Rosenblatt.
Rigidity and non-recurrence along sequences.
Ergodic Theory Dynam. Systems, 34(5):1464–1502, 2014.
- [6] V. Bergelson and A. Leibman.
Polynomial extensions of van der Waerden's and Szemerédi's theorems.
J. Amer. Math. Soc., 9(3):725–753, 1996.

- [7] V. Bergelson and A. Leibman.
Failure of the Roth theorem for solvable groups of exponential growth.
Ergod. Th. and Dynam. Sys., 24:45–53, 2004.
- [8] A. I. Bufetov and B. Solomyak.
On substitution automorphisms with pure singular spectrum.
Math. Z., 301(2):1315–1331, 2022.
- [9] M. Einsiedler and T. Ward.
Ergodic theory with a view towards number theory, volume 259 of *Graduate Texts in Mathematics*.
Springer-Verlag London, Ltd., London, 2011.

References IV

- [10] S. Farhangi.
Topics in ergodic theory and ramsey theory.
PhD dissertation, the Ohio State University, 2022.
- [11] B. Fayad.
Smooth mixing flows with purely singular spectra.
Duke Math. J., 132(2):371–391, 2006.
- [12] N. Frantzikinakis.
Furstenberg systems of Hardy field sequences and applications.
J. Anal. Math., 147(1):333–372, 2022.
- [13] N. Frantzikinakis and B. Host.
Multiple recurrence and convergence without commutativity.
arXiv:2111.01518v2 [math.DS], 2021.

- [14] N. Frantzikinakis, E. Lesigne, and M. Wierdl.
Sets of k -recurrence but not $(k + 1)$ -recurrence.
Ann. Inst. Fourier (Grenoble), 56(4):839–849, 2006.
- [15] N. Frantzikinakis, E. Lesigne, and M. Wierdl.
Random sequences and pointwise convergence of multiple ergodic averages.
Indiana Univ. Math. J., 61(2):585–617, 2012.
- [16] K. Frączek and M. Lemańczyk.
On mild mixing of special flows over irrational rotations under piecewise smooth functions.
Ergodic Theory Dynam. Systems, 26(3):719–738, 2006.

References VI

[17] H. Furstenberg.

Ergodic behavior of diagonal measures and a theorem of Szemerédi on arithmetic progressions.

J. Analyse Math., 31:204–256, 1977.

[18] H. Furstenberg.

Recurrence in ergodic theory and combinatorial number theory.

Princeton University Press, Princeton, N.J., 1981.

M. B. Porter Lectures.

[19] B. M. Gurevič.

The entropy of horocycle flows.

Dokl. Akad. Nauk SSSR, 136:768–770, 1961.

References VII

[20] B. Host.

Mixing of all orders and pairwise independent joinings of systems with singular spectrum.

Israel J. Math., 76(3):289–298, 1991.

[21] I. Klemes and K. Reinhold.

Rank one transformations with singular spectral type.

Israel J. Math., 98:1–14, 1997.

[22] O. S. Parasyuk.

Flows of horocycles on surfaces of constant negative curvature.

Uspehi Matem. Nauk (N.S.), 8(3(55)):125–126, 1953.

References VIII

[23] V. A. Rohlin.

Lectures on the entropy theory of transformations with invariant measure.

Uspehi Mat. Nauk, 22(5 (137)):3–56, 1967.

[24] S. V. Tikhonov.

A complete metric on the set of mixing transformations.

Mat. Sb., 198(4):135–158, 2007.

[25] A. Tserunyan.

A Ramsey theorem on semigroups and a general van der Corput lemma.

J. Symb. Log., 81(2):718–741, 2016.

[26] J. G. van der Corput.

Diophantische Ungleichungen. I. Zur Gleichverteilung Modulo Eins.

Acta Math., 56(1):373–456, 1931.